

# Milk Consumption and Prostate Cancer

**By Neal D. Barnard, M.D.**

## **Abstract**

Prostate cancer is one of the most common malignancies worldwide, with an estimated 400,000 new cases diagnosed annually. Its incidence and mortality have been associated with milk or dairy product consumption in international and interregional correlational studies. As a result, case-control and cohort studies have further investigated this association and are described in this review. Of 12 case-control studies, six found significant associations, as did five of 11 cohort studies, with relative risk of prostate cancer among those with the most frequent dairy product consumption ranging between 1.3 and 2.5, with evidence of a dose-response relationship. Mechanisms that may explain this association include the deleterious effect of high-calcium foods on vitamin D balance, the tendency of frequent dairy intake to increase serum insulin-like growth factor I (IGF-1) concentrations, and the effect of dairy products on testosterone concentration or activity.

## **Introduction**

Prostate cancer is the fourth most common malignancy among men worldwide, with an estimated 400,000 new cases diagnosed annually, accounting for 3.9 percent of all new cancer cases.<sup>1</sup> Epidemiologic evidence strongly suggests that dietary factors play a major role in prostate cancer progression and mortality, with protective effects associated with consumption of fruit (particularly tomatoes), vitamin E, and selenium, and increased risk linked to dairy products, meat, and fat.<sup>2</sup> Dairy product consumption has been associated with prostate cancer risk in divergent populations, and several studies have investigated mechanisms that may explain these findings. This review describes studies reporting prostate cancer risk in relation to milk or dairy products generally. It does not include studies reporting only intake of individual milk derivatives, such as butter, or nutrients, such as fat.

## **Correlational Studies**

In international and interregional correlational studies, dairy product consumption has been consistently associated with prostate cancer mortality.<sup>3-7</sup> The largest and most recent of these, based on World Health Organization mortality figures for 1985-1989 from 59 countries and United Nations food balance data for 1979-1981, reported a strong correlation between per capita milk consumption and prostate cancer mortality ( $r = 0.78$ ,  $P < 0.0001$ ).<sup>7</sup> A more geographically restricted study, conducted in 20 Italian regions, found a similar correlation between prostate cancer mortality and milk consumption ( $r = 0.75$ ,  $P < 0.01$ ).<sup>6</sup>

International correlational studies typically rely on food "disappearance" data, which may not accurately reflect intake, and are limited in their ability to control for potential confounders. They are also subject to variability in reporting practices, although this is less likely to influence mortality data than incidence figures. Some of the weaknesses of international correlational studies are avoided in case-control and cohort studies. Case-control studies compare the recalled diets of individuals with cancer with the diets of individuals without cancer who are similar in other relevant respects. Because cases and locally recruited controls are likely to have similar dietary patterns, the sensitivity of such studies is often limited. Cohort studies gather dietary information from healthy volunteers who are then followed over time.

## **Case-Control Studies**

Six case-control studies in geographically diverse areas have reported significantly increased risk of prostate cancer (relative risk ranging from 1.5 to 2.5) for those in the highest categories of dairy product consumption, compared with lower consumption levels ([Table 1](#)).<sup>8-13</sup> Four additional studies reported nonsignificant positive associations, and two found no association between dairy product consumption and prostate cancer incidence.<sup>14-19</sup>

Two studies in northern Italy compared prostate cancer patients with hospital controls, finding increased risk of prostate cancer among those with the most frequent milk consumption.<sup>8-10</sup> Similarly, a study at Roswell Park Memorial Institute

in Buffalo, N.Y., found an increased risk of prostate cancer with the daily consumption of three or more glasses of whole milk, compared with never drinking milk.<sup>9</sup>In Montevideo, Uruguay, a comparison of prostate cancer patients with hospital controls, most of whom had other forms of cancer, found an increased risk of prostate cancer associated with drinking two or more milk servings per day, compared with having less than one serving per day.<sup>11</sup> In Örebro county, Sweden, men with prostate cancer were compared with controls selected from a population register. Higher dairy product consumption was associated with increased relative risk of prostate cancer.<sup>12</sup>A preliminary study from a portion of this patient sample found no associations between prostate cancer and any food recalled as having been consumed during adolescence.<sup>20</sup>A U.S. study compared men newly diagnosed with prostate cancer and healthy population-based controls in Georgia, New Jersey, and Michigan. Dairy product consumption was associated with prostate cancer risk among whites, but not blacks.<sup>13</sup>

These studies have the methodologic strengths of statistical adjustment for age and other factors and reasonably large sample sizes ([Table 1](#)). Of those studies finding positive but nonsignificant associations between dairy use and prostate cancer, several used smaller sample sizes or failed to adjust for age or other variables ([Table 1](#)). In Los Angeles and Chicago, prostate cancer patients were matched to hospital controls, finding nonsignificant increases in dairy product consumption among cancer patients.<sup>14</sup> A Minnesota study comparing prostate cancer patients with hospital and neighborhood controls reported nonsignificant increases in dairy product consumption among cancer patients.<sup>15</sup> Similarly, a small study in Japan comparing prostate cancer patients to healthy controls from a prostate cancer-screening program found a nonsignificant increased risk associated with daily milk consumption.<sup>16</sup> In Athens, Greece, prostate cancer patients were compared with hospital controls. Milk and dairy product consumption was marginally positively associated with prostate cancer risk.<sup>17</sup>

A Swedish case-control study including men with prostate cancer and unrelated controls drawn from a twin registry found no relationship between cancer risk and any dietary factor.<sup>18</sup> An English study compared prostate cancer patients to controls

with benign prostatic hyperplasia (BPH) and hospital controls with non-urological disease; data for both groups of controls were combined. The study reported no association between dairy product consumption and prostate cancer risk. The use of BPH patients as controls in this study may have reduced its sensitivity, as BPH may have antecedents similar to those of prostate cancer.<sup>19</sup>

## **Cohort Studies**

Five of 11 cohort studies have found significant associations between milk or dairy product consumption and prostate cancer incidence or mortality.<sup>21-25</sup> Six studies found no association between milk or dairy product use generally and prostate cancer incidence or mortality.<sup>26-31</sup>

A 20-year study of prostate cancer mortality among California Seventh-day Adventists reported a dose-related increased risk of age-adjusted prostate cancer mortality with milk consumption (for  $\geq 3$  glasses daily, RR = 2.4, 95% CI, 1.3-4.3; for 1-2 glasses daily, RR = 1.8, 95% CI, 1.0-3.0, compared to  $< 1$  glass per day.) In a multivariate analysis adjusting for age, education, body weight, and consumption of meat, milk, cheese, and eggs, the relative risk of fatal prostate cancer associated with drinking  $\geq 3$  glasses of milk per day was reduced to 1.5 and was no longer statistically significant ( $p < 0.10$ ).<sup>21</sup> However, adjustment for cheese consumption may be inappropriate if the relevant dietary factor is dairy product consumption generally. Similarly, adjustment for body weight may be inappropriate if increased body weight is one of the mechanisms by which dairy product consumption influences prostate cancer risk. A separate study of California Adventists studied cancer incidence, rather than mortality, finding no relationship with milk consumption.<sup>27</sup>

A dose-response relationship was also suggested by a cohort study including various ethnicities in Hawaii. Relative risks of prostate cancer, adjusted for age, ethnicity, and income, for men in the middle and highest tertiles of milk consumption were 1.3 (CI, 1.0-1.9) and 1.4 (CI, 1.0-2.1), respectively, compared with the lowest tertile. Although these 95 percent confidence intervals included 1.0, a statistically significant trend was reported ( $P_{\text{trend}} = 0.04$ ).<sup>22</sup>

In the Health Professionals Follow-Up Study, a cohort of U.S. male dentists, optometrists, osteopaths, pharmacists, and veterinarians, relative risk of advanced prostate cancer associated with daily consumption of more than two glasses of milk, compared to zero, was 1.6 (95% CI, 1.2-2.1,  $P_{\text{trend}} = 0.002$ ). For metastatic disease, relative risk was 1.8 (95% CI, 1.2-2.8,  $P_{\text{trend}} = 0.01$ ). Of the milk consumed, 83 percent was skim or low-fat.<sup>23</sup>

The Netherlands Cohort Study reported a trend of increased prostate cancer risk with increasing milk consumption after adjustment for age, family history of prostate cancer, and socioeconomic status, although the difference in risk, compared with the lowest (index) quintile of milk consumption, was significant only for the 4th quintile (RR= 1.63, 95% CI, 1.20-2.20,  $P_{\text{trend}} = 0.02$ ).<sup>24</sup>

In the Physicians' Health Study cohort, consumption of two and one-half dairy servings daily was associated with increased risk of prostate cancer, compared with having less than one-half serving daily, after adjustment for age, smoking, exercise level, and body mass index (BMI).<sup>25</sup>

Among the studies finding no association between dairy product use and prostate cancer, one, conducted in Norway, found a significantly increased age-adjusted risk of prostate cancer with consumption of skim milk, compared with whole milk (incidence rate ratio 2.2, 95% CI, 1.3-3.7), although milk consumption in general was not associated with risk. The authors speculate that the relatively young age of their sample (mean age 43 years, range 16-56 years, at the outset of a nine- to 15-year follow-up period) may have reduced study sensitivity.<sup>31</sup>

Hirayama found a protective effect of green and yellow vegetables, but no detectable effect of milk consumption. Although the cohort was large (112,261 men), it identified only 63 cancer deaths during the follow-up period, and did not limit the inclusion of the oldest participants. Milk consumption was probably uncommon in this group, but the number of men consuming milk with various frequencies was not reported.<sup>26</sup> In a cohort of men of Japanese ancestry living in Hawaii, there was no association between milk consumption and age-adjusted prostate cancer risk. Milk consumption was uncommon; only 34 percent of cases

consumed milk five times per week or more.<sup>28</sup> In a Rancho Bernardo, Calif., cohort (aged 50-84 at the study's outset), no relationship was found between whole milk consumption and prostate cancer incidence during 14-year follow-up, after adjustment for age, history of heart disease or diabetes, BMI, systolic blood pressure, smoking, and plasma cholesterol concentration. Milk consumption averaged 0.5 cups per day.<sup>29</sup> In a cohort of white male policyholders of the Lutheran Brotherhood Insurance Society, most of whom lived in Minnesota and the northeastern U.S., no associations were identified between prostate cancer mortality and any dietary factor. The authors cautioned that the lack of an association between dietary factors and cancer risk may be partially due to the limited number of items in the food-frequency questionnaire and the homogeneous nature of the cohort, heavily weighted toward individuals of Scandinavian descent.<sup>30</sup>

In summary, six of 12 case-control studies and five of 11 cohort studies found significant associations between milk or dairy product consumption and prostate cancer incidence and mortality. Particularly among cohort studies, those reporting significant associations were generally larger and more recent.

## **Mechanisms**

Dairy products may influence the incidence or progression of prostate cancer by several possible mechanisms. One for which evidence is particularly compelling is the effect of high-calcium foods on vitamin D metabolism. In several prospective studies, calcium intake has emerged as an independent predictor of prostate cancer risk.<sup>12,23,25</sup> Vitamin D is derived either by conversion from 7-dehydrocholesterol in a reaction catalyzed by ultraviolet light, or from dietary sources. For conversion to the biologically active hormone, a hydroxyl group is added in the liver to form 25(OH) vitamin D, and a second hydroxyl group is added in the kidney, producing 1,25(OH)<sub>2</sub> vitamin D.

Vitamin D receptors are present on prostate epithelium. Among other functions, 1,25(OH)<sub>2</sub>vitamin D reduces cell proliferation and enhances cell differentiation.<sup>32</sup> An oral calcium load suppresses parathyroid hormone secretion which, in turn, reduces renal 1,25(OH)<sub>2</sub> vitamin D production. Calcium itself also downregulates this

reaction. Although some dairy products are supplemented with vitamin D, this inactive prehormone does not appreciably increase 1,25(OH)<sub>2</sub> vitamin D blood levels, and the net effect of dairy consumption is a reduction in blood levels of this active form of the hormone. Milk's high content phosphorus and animal protein may aggravate this effect.<sup>32</sup>

Dairy product consumption has also been shown to increase serum concentrations of insulin-like growth factor (IGF-1).<sup>33,34</sup> In in-vitro studies, IGF-1 has mitogenic and antiapoptotic properties on prostate epithelial cells.<sup>35,36</sup> Case-control studies in diverse populations have shown a strong and consistent association between serum IGF-1 concentrations and prostate cancer risk.<sup>37</sup> In men and women aged 55 to 85 years, the addition of three daily eight-ounce servings of nonfat or 1 percent milk for 12 weeks was associated with a 10 percent increase in serum IGF-1 concentration ( $P < 0.001$ ).<sup>34</sup> Mean serum IGF-1 concentration among vegans was shown to be 8 percent lower than among ovolactovegetarians ( $P = 0.01$ ) and 9 percent lower than among meat-eaters ( $P = 0.01$ ).<sup>38</sup> Changing dietary protein sources from animal sources to plant sources has been shown to reduce serum IGF-1 concentrations.<sup>39</sup>

Most dairy products contain substantial amounts of fat and are devoid of fiber, a combination that is likely to increase serum testosterone concentration and activity, with a mitogenic effect on prostate tissue.<sup>40</sup> However, several studies have found an association of dairy product intake with prostate cancer incidence and mortality that is independent of total fat intake and other dietary variables.<sup>25,40,41</sup>

Dietary factors other than dairy products are also associated with risk. Generally speaking, diets high in animal products are associated with higher risk, while those rich in plant foods, particularly tomatoes, are associated with reduced risk.<sup>2</sup> Daily soymilk consumption was associated with a significant reduction in prostate cancer risk in a cohort of 13,855 Seventh-Day Adventist men (RR = 0.3, 95% CI, 0.1-1.0, compared with those never drinking soymilk).<sup>42</sup> Isoflavones in soymilk inhibit growth of human prostate cancer cells and also inhibit 5 $\alpha$ -reductase, an enzyme that converts testosterone to 5 $\alpha$ -dihydrotestosterone in the prostate.<sup>42</sup> A similar beneficial effect was demonstrated for tofu consumption.<sup>28</sup> Based on experience

with a case-control study in Athens, Greece, researchers calculated that the combined effect of reducing dairy consumption, substituting olive oil for other added fats, and increasing tomato intake to the levels consumed by those in the lowest risk categories could reduce prostate cancer risk in their population by 41 percent (95% CI, 23-59%).<sup>43</sup>

## **Conclusions**

Evidence from international, case-control, and cohort studies suggests that men who avoid dairy products are at lower risk for prostate cancer incidence and mortality, compared with others. In case-control and cohort studies, the relative risk of prostate cancer among subgroups with the most frequent milk consumption, compared with those at the lowest consumption levels, falls in the range of 1.3 to 2.5. These findings raise two important questions: Does the observed relationship represent cause and effect, and is available evidence sufficient to justify a recommendation that milk-drinking men alter their dietary habits?

Findings supporting a cause-and-effect relationship include the relative consistency of this association in diverse populations, evidence of a dose-response relationship, plausible biological mechanisms that underlie the observed associations, and no reasonable alternative explanation for these findings. Perspective is lent to the second question by a comparison with evidence linking alcohol use and breast cancer risk. Although somewhat fewer studies have addressed the association between milk and prostate cancer, their demonstrated effect, strength, and consistency of evidence approach those relating alcohol to breast cancer risk, an association that is now widely accepted and incorporated into the Dietary Guidelines for Americans.<sup>44</sup> A pooled analysis of cohort studies showed that the adjusted relative risk of incident breast cancer for women consuming two to five drinks (30-60 grams of alcohol) per day was 1.41 (95% CI, 1.18-1.69).<sup>45</sup> In a 1997 review by the World Cancer Research Fund and the American Institute for Cancer Research, six of 11 cohort studies and 15 of 36 case-control studies found such an association.<sup>46</sup>



Men who choose to avoid dairy products reap other nutritional benefits, such as a reduction in total fat, saturated fat, and cholesterol intake. Unless they replace dairy products with calcium-fortified products or calcium supplements, they are likely to reduce their calcium intake in the process. However, a reduction in calcium intake may be an important mechanism by which reducing or avoiding dairy products reduces prostate cancer risk. Moreover, there is no apparent risk to moderate reductions in calcium intake. There is little evidence to suggest that a high intake of calcium from dairy or other sources reduces the risk of osteoporotic fractures among men. Few studies have examined the effect of dietary calcium on osteoporosis risk in adult men independently of vitamin D intake.<sup>47</sup> There has been some indication that a higher intake of calcium, including that from dairy sources, in the context of an omnivorous American diet, is associated with reduced recurrence of colonic adenomatous polyps.<sup>48</sup> However, in Africa, in the context of a low-calcium, low-dairy diet, both adenomatous polyps and colon cancer are much rarer than in Western countries.<sup>49</sup> Some studies suggest that calcium, including that in dairy products, may reduce blood pressure, but the effect, if any, is small (on the order of <2 mm Hg systolic and <1 mm Hg diastolic), far smaller than the effect of adding vegetables and fruits to the diet.<sup>50-51</sup>

In conclusion, several lines of evidence indicate that consumption of dairy products is associated with increased risk of prostate cancer incidence and mortality. Avoidance of these products may offer a means of reducing risk of this common illness.

## References

1. World Cancer Research Fund/American Institute for Cancer Research. Food, Nutrition, and the Prevention of Cancer: A Global Perspective. American Institute for Cancer Research, Washington, D.C., 1997, p. 311.
2. Chan JM, Stampfer MJ, Giovannucci EL. What causes prostate cancer? A brief summary of the epidemiology. *Sem Canc Biol.* 1998a;8:263-73.
3. Howell MA. Factor analysis of international cancer mortality data and per capita food consumption. *Br J Cancer.* 1974;29:328-36.
4. Armstrong B, Doll R. Environmental factors and cancer incidence and mortality in different

- countries, with special reference to dietary practices. *Int J Cancer*. 1975;15: 617-31.
5. Rose DP, Boyar AP, Wynder EL. International comparisons of mortality rates for cancer of the breast, ovary, prostate, and colon, and per capita food consumption. *Cancer*. 1986;58:2363-71.
  6. Decarli A, La Vecchia C. Environmental factors and cancer mortality in Italy: correlational exercise. *Oncology*. 1986;43:116-26.
  7. Hebert JR, Hurley TG, Olendzki BC, Teas J, Ma Y, Hampl JS. Nutritional and socioeconomic factors in relation to prostate cancer mortality: a cross national study. *J Natl Cancer Inst*. 1998;90(21): 1637-47.
  8. Talamini R, La Vecchia C, Decarli A, Negri E, Franceschi S. Nutrition, social factors and prostatic cancer in a Northern Italian population. *Br J Cancer*. 1986;53:817-21.
  9. Mettlin C, Selenskas S, Natarajan N, Huben R. Beta-carotene and animal fats and their relationship to prostate cancer risk. *Cancer*. 1989;64:605-12.
  10. Talamini R, Franceschi S, La Vecchia C, Serraino D, Barra S, Negri E. Diet and prostate cancer: a case-control study in Northern Italy. *Nutr Cancer*. 1992;118:277-86.
  11. De Stefani E, Fierro L, Barrios E, Ronco A. Tobacco, alcohol, diet and risk of prostate cancer. *Tumori*. 1995;81:315-20.
  12. Chan JM, Giovannucci EL, Andersson SO, Yuen J, Adami HO, Wolk A. Dairy products, calcium, phosphorous, vitamin D, and risk of prostate cancer (Sweden). *Cancer Causes and Control*. 1998b; 9:559-66.
  13. Hayes RB, Ziegler RG, Gridley G, Swanson C, Greenberg RS, Swanson GM, Schoenberg JB, Silverman DT, Brown LM, Pottern LM, Liff J, Schwartz AG, Fraumeni JF Jr, Hoover RN. Dietary factors and risks for prostate cancer among blacks and whites in the United States. *Cancer Epidemiol Biomarkers Prev*. 1999;8:25-34.
  14. Rotkin ID. Studies in the epidemiology of prostatic cancer: expanded sampling. *Cancer Treatment Reports*. 1977;61:173-80.
  15. Schuman LM, Mandel JS, Radke A, Seal U, Halberg F. Some selected features of the epidemiology of prostatic cancer: Minneapolis-St. Paul, Minnesota case-control study, 1976-1979. In Magnus K, Ed. *Trends in Cancer Incidence: Causes and Practical Implications*, Hemisphere Publishing, Washington, D.C.; 1982:345-54.
  16. Mishina T, Watanabe H, Araki H, Nakao M. Epidemiological study of prostatic cancer by matched-pair analysis. *Prostate*. 1985;6:423-36.
  17. Tzonou A, Signorello LB, Lagiou P, Wu J, Trichopoulos D, Trichopoulou A. Diet and cancer of the

prostate: a case-control study in Greece. *Int J Cancer*. 1999;80:704-8.

18. Grönberg H, Damber L, Damber JE. Total food consumption and body mass index in relation to prostate cancer risk: a case-control study in Sweden with prospectively collected exposure data. *J Urology*. 1996;155:969-74.

19. Ewings P, Bowie C. A case-control study of cancer of the prostate in Somerset and east Devon. *Br J Cancer*. 1996;74:661-6.

20. Andersson SO, Baron J, Wolk A, Lindgren C, Bergstrom R, Adami HO. Early life risk factors for prostate cancer: a population-based case-control study in Sweden. *Cancer Epidemiol Biomark Prev*. 1995;4:187-92.

21. Snowdon DA, Phillips RL, Choi W. Diet, obesity, and risk of fatal prostate cancer. *Am J Epidemiology*. 1984;120:244-50.

22. LeMarchand L, Kolonel LN, Wilkens LR, Myers BC, Hirohata T. Animal fat consumption and prostate cancer: a prospective study in Hawaii. *Epidemiology*. 1994;5:276-82.

23. Giovannucci E, Rimm EB, Wolk A, Ascherio A, Stampfer MJ, Colditz GA, Willett WC. Calcium and fructose intake in relation to risk of prostate cancer. *Cancer Res*. 1998a;58:442-7.

24. Schuurman AG, van den Brandt PA, Dorant E, Goldbohm RA. Animal products, calcium and protein and prostate cancer risk in the Netherlands Cohort Study. *Br J Cancer*. 1999;80:1107-1113.

25. Chan JM, Stampfer MJ, Ma J, Gann PH, Gaziano JM, Giovannucci E. Dairy products, calcium, and prostate cancer risk in the Physicians' Health Study. *Am J Clin Nutr*. 2001;74:549-54.

26. Hirayama T. Epidemiology of prostate cancer with special reference to the role of diet. *Natl Cancer Inst Monogr*. 1979;53:149-55.

27. Mills PK, Beeson WL, Phillips RL, Fraser GE. Cohort study of diet, lifestyle, and prostate cancer in Adventist men. *Cancer*. 1989;64:598-604.

28. Severson RK, Nomura AMY, Grove JS, Stemmermann GN. A prospective study of demographics, diet, and prostate cancer among men of Japanese ancestry in Hawaii. *Cancer Res*. 1989;49:1857-60.

29. Thompson MM, Garland C, Barrett-Connor E, Khaw KT, Friedlander NJ, Wingard DL. Heart disease risk factors, diabetes, and prostatic cancer in an adult community. *Am J Epidemiol*. 1989;129:511-7.

30. Hsing AW, McLaughlin JK, Schuman LM, Bjelke E, Gridley G, Wacholder S, Co Chien HT, Blot WJ. Diet, tobacco use, and fatal prostate cancer: results from the Lutheran brotherhood cohort study. *Cancer Res*. 1990;50:6836-40.

31. Veierød MB, Laake P, Thelle DS. Dietary fat intake and risk of prostate cancer: a prospective study of 25,708 Norwegian men. *Int J Cancer*. 1997;73:634-8.

32. Giovannucci E. Dietary influences of 1,25(OH)<sub>2</sub> vitamin D in relation to prostate cancer: a hypothesis. *Cancer Causes and Control*. 1998b;9:567-82.
33. Cadogan J, Eastell R, Jones N, Barker ME. Milk intake and bone mineral acquisition in adolescent girls: randomised, controlled intervention trial. *BMJ*. 1997;315:1255-60.
34. Heaney RP, McCarron DA, Dawson-Hughes B, Oparil S, Berga SL, Stern JS, Barr SI, Rosen CJ. Dietary changes favorably affect bone remodeling in older adults. *J Am Dietetic Asso*. 1999;99:1228-33.
35. Iwamura M, Sluss PM, Casamento JB, Cockett ATK. Insulin-like growth factor I: action and receptor characterization in human prostate cancer cell lines. *Prostate*. 1993;22:243-52.
36. Culig Z, Hobisch A, Cronauer MV, Radmayr C, Trapman J, Hittmair A, Bartsch G, Klocker H. Androgen receptor activation in prostatic tumor cell lines by insulin-like growth factor-I, keratinocyte growth factor, and epidermal growth factor. *Eur Urol*. 1995;27(suppl 2):45-7.
37. Cohen P. Serum insulin-like growth factor-I levels and prostate cancer risk—interpreting the evidence. *J Natl Cancer Inst*. 1998;90:876-9.
38. Allen NE, Appleby PN, Davey GK, Key TJ. Hormones and diet: low insulin-like growth factor-I but normal bioavailable androgens in vegan men. *Br J Cancer*. 2000;83:95-7.
39. Kontessis PA, Bossinakou I, Sarika L, Iliopoulou E, Papantoniou A, Trevisan R, Roussi D, Stipsanelli K, Grigorakis S, Souvatzoglou A. Renal, metabolic, and hormonal responses to proteins of different origin in normotensive, nonproteinuric type I diabetic patients. *Diabetes Care* 1995;18:1233-40.
40. Ross RK, Henderson BE. Do diet and androgens alter prostate cancer risk via a common etiologic pathway? *J Natl Cancer Inst*. 1994;86:252-4.
41. Giovannucci E, Rimm EB, Colditz GA, Stampfer MJ, Ascherio A, Chute CC, Willett WC. A prospective study of dietary fat and risk of prostate cancer. *J Natl Cancer Inst*. 1993;85:1571-9.
42. Jacobsen BK, Knutsen SF, Fraser GE. Does high soy milk intake reduce prostate cancer incidence? The Adventist Health Study (United States). *Cancer Causes and Control*. 1998;9:553-7.
43. Bosetti C, Tzonou A, Lagiou P, Negri E, Trichopoulos D, Hsieh C-C. Fraction of prostate cancer incidence attributed to diet in Athens, Greece. *Eur J Cancer Prev*. 2000;9:119-23.
44. U.S. Department of Agriculture and Department of Health and Human Services. Nutrition and Your Health: Dietary Guidelines for Americans, 5th Edition. Home and Garden Bulletin No. 232. U.S. Department of Agriculture, Washington, D.C., 2000, p. 36.
45. Smith-Warner SA, Spiegelman D, Yuan SS, van den Brandt PA, Folsom AR, Goldbohm A, Graham S, Holmberg L, Howe GR, Marshall JR, Miller AB, Potter JD, Speizer FE, Willett WC, Wolk A, Hunter DJ.

Alcohol and breast cancer among women: a pooled analysis of cohort studies. *JAMA*.

1998;279:535-40.

46. World Cancer Research Fund/American Institute for Cancer Research. Food, Nutrition, and the Prevention of Cancer: A Global Perspective. American Institute for Cancer Research, Washington, D.C., 1997, pp. 270-74.

47. Welten DC, Kemper HCG, Post GB, Van Staveren WA. A meta-analysis of the effect of calcium intake on bone mass in young and middle-aged females and males. *J Nutr*. 1995;125:2802-13.

48. Holt PR, Attilsoy EO, Gilman J, Guss J, Moss SF, Newmark H, Fan K, Yang K, Lipkin M. Modulation of abnormal colonic epithelial cell proliferation and differentiation by low-fat dairy foods: a randomized controlled trial. *JAMA*. 1998;280:1074-9.

49. Walker ARP, Segal I. Low-fat dairy foods and colonic epithelial cell proliferation (letter). *JAMA*. 1999;281:1274.

50. Griffith LE, Guyatt GH, Cook RJ, Bucher HC, Cook DJ. The influence of dietary and nondietary calcium supplementation on blood pressure: an updated metaanalysis of randomized controlled trials. *Am J Hypertension*. 1999;12:84-92.

51. Sacks FM, Appel LJ, Moore TJ, Obarzanek E, Vollmer WM, Svetkey LP, Bray GA, Vogt TM, Cutler JA, Windhauser MM, Lin PH, Karanja N. A dietary approach to prevent hypertension: a review of the Dietary Approaches to Stop Hypertension (DASH) study. *Clin Cardiol*. 1999;22(suppl III):III-6-III-10.